

A Deduced Feynman Rule for Calculating Retarded and Advanced Green Function in Closed Time Path Formalism

Jun Xiao and Enke Wang

Institute of Particle Physics, Huazhong Normal University, Wuhan 430079

(February 1, 2008)

Based on the closed time path formalism, a new Feynman rule for directly calculating the retarded and advanced Green functions is deduced. This Feynman rule is used to calculate the two-point self-energy and three-point vertex correction in ϕ^3 theory. The generalized fluctuation-dissipation theorem for three-point nonlinear response function is verified.

The closed time path formalism (CTPF) [1–3] in real-time finite-temperature field theory has been used widely to investigate the equilibrium and non-equilibrium properties [4–9] of the thermal system recently. It is different to the imaginary time formalism (ITF), without analytical continuation the physical region can be reached in the real time formalism. In the CTPF two kinds of fields are defined according to the closed time integral contour in the generating function of Green function. The closed time integral contour consists of the first branch which runs from negative infinity to positive infinity and the second branch which runs back from positive infinity to negative infinity. The field located on the first and the second branch is defined as physical (type-1) and ghost (type-2) field, respectively. So that the propagator becomes a 2×2 matrix with components $G_{11}, G_{12}, G_{21}, G_{22}$, their corresponding self-energy are denoted as $\Sigma_{11}, \Sigma_{12}, \Sigma_{21}, \Sigma_{22}$ according to the type-1 or type-2 external leg of the Feynman diagram. It is known that the self-energy with physical interpretation is not Σ_{11} but the retarded function Σ^{ret} (or advanced function Σ^{adv}) which is the analytical continuation of the self-energy obtained in the ITF [9]. In order to calculate retarded (or advanced) self-energy in the real time formalism, the usual way is to calculate $\Sigma_{11}, \Sigma_{12}, \Sigma_{21}$ and Σ_{22} first, and then use the relation $\Sigma^{ret} = \Sigma_{11} + \Sigma_{12}$ (or $\Sigma^{adv} = \Sigma_{11} + \Sigma_{21}$) to get them. When one calculates n -point ($n > 2$) retarded/advanced vertex [10], for example, three-point fully retarded vertex Γ_{raa} defined in Ref. [11], we need to calculate eight components $\Gamma_{111}, \Gamma_{112}, \Gamma_{121}, \Gamma_{122}, \Gamma_{211}, \Gamma_{212}, \Gamma_{221}$ and Γ_{222} first and then use the relation $\Gamma_{raa} = \frac{1}{2}(\Gamma_{111} - \Gamma_{112} - \Gamma_{121} + \Gamma_{122} + \Gamma_{211} - \Gamma_{212} - \Gamma_{221} + \Gamma_{222})$ to get it, the calculation becomes somewhat tedious. In this letter we deduce a novel Feynman rule which can be used to calculate n -point retarded/advanced Green function directly to avoid above tedious calculation, and then we verify the generalized Fluctuation-dissipation theorem for three-point nonlinear response function by calculating the vertex correction.

In the CTPF, for any field ϕ the four components of the matrix propagator in the single-time representation are defined as

$$G_{a_1 a_2}(x_1, x_2) \equiv -i \langle T_p(\phi_{a_1}(x_1) \phi_{a_2}(x_2)) \rangle, \quad (1)$$

where T_p represents the time ordering operator along the closed time path, it is normal and antichronological time ordering on the first and second branch of the closed time path, respectively. $a_1, a_2 \in \{1, 2\}$ indicate on which of the two branches the ϕ fields are located. $\langle \cdots \rangle$ stands for the thermal expectation value. Following [3] we define

$$\phi_a(x) = \phi_1(x) - \phi_2(x), \quad \phi_r(x) = \frac{1}{2}(\phi_1(x) + \phi_2(x)), \quad (2)$$

and the 2-point Green function in the above (r, a) basis

$$G_{\alpha_1 \alpha_2}(x_1, x_2) \equiv -i 2^{n_r-1} \langle T_p(\phi_{\alpha_1}(x_1) \phi_{\alpha_2}(x_2)) \rangle, \quad (3)$$

where $\alpha_1, \alpha_2 \in \{a, r\}$, and n_r is the number of r indices among $\{\alpha_1, \alpha_2\}$. Inserting Eq. (2) into (3) it is easy to get

$$G_{rr}(x_1, x_2) = G_{12}(x_1, x_2) + G_{21}(x_1, x_2) \\ = -i \langle [\phi(x_1), \phi(x_2)]_{\pm} \rangle, \quad (4)$$

$$G_{ra}(x_1, x_2) = G_{11}(x_1, x_2) - G_{12}(x_1, x_2) \\ = -i \theta(x_1^0 - x_2^0) \langle [\phi(x_1), \phi(x_2)]_{\mp} \rangle, \quad (5)$$

$$G_{ar}(x_1, x_2) = G_{11}(x_1, x_2) - G_{21}(x_1, x_2) \\ = i \theta(x_2^0 - x_1^0) \langle [\phi(x_1), \phi(x_2)]_{\mp} \rangle, \quad (6)$$

$$G_{aa}(x_1, x_2) = 0. \quad (7)$$

Here $[\phi(x_1), \phi(x_2)]_{\pm} = \phi(x_1)\phi(x_2) \pm \phi(x_2)\phi(x_1)$. The double sign $\pm$ or $\mp$ in Eq. (4)-(6) corresponds to Boson and Fermion field for the upper and lower case, respectively. Obviously $G_{ra}(x_1, x_2)$ and $G_{ar}(x_1, x_2)$ are the usual retarded and advanced Green functions.

Denote the 2×2 matrix propagator in the single-time representation as

$$G = \begin{pmatrix} G_{11} & G_{12} \\ G_{21} & G_{22} \end{pmatrix}, \quad (8)$$

and the matrix propagator in the (r, a) representation as

$$\overline{G} = \begin{pmatrix} G_{aa} & G_{ar} \\ G_{ra} & G_{rr} \end{pmatrix} = \begin{pmatrix} 0 & G^{adv} \\ D^{ret} & D^{col} \end{pmatrix}. \quad (9)$$

Both matrix propagators satisfy following transformation relation [3]:

$$G = Q^\dagger \bar{G} Q, \quad (10)$$

where Q is orthogonal Keldysh transformation matrix

$$Q = \begin{pmatrix} Q_{a1} & Q_{a2} \\ Q_{r1} & Q_{r2} \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}, \quad (11)$$

$$Q^\dagger = \begin{pmatrix} Q_{1a}^\dagger & Q_{1r}^\dagger \\ Q_{2a}^\dagger & Q_{2r}^\dagger \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}. \quad (12)$$

From the above transformation (10) we see that, both left and right side of the propagator $\bar{G}$ associate with the transformation matrix Q . We can establish a new Feynman rule in following way: as illustrated in Fig. 1, the matrices $Q^\dagger$ can be absorbed into the left vertex with outgoing momentum line, and Q can be absorbed into the right vertex with incoming momentum line, so that we leave $\bar{G}_{\alpha\beta}$ in the (r, a) basis as a new propagator of the propagating line. After absorbing all Q from propagating line, the new bare vertex with all incoming momentum (as illustrated in Fig. 2(a)) can be defined as

$$\gamma_{\beta\beta'\dots\beta''}(p, q, \dots r) = Q_{\beta a} Q_{\beta' b} \dots Q_{\beta'' c} g_{ab\dots c}, \quad (13)$$

where $a, b, \dots, c \in 1, 2$ and $\beta, \beta', \dots, \beta'' \in a, r$, a summation over repeated indices is understood, $p + q + \dots + r = 0$ because of energy-momentum conservation. In the single-time representation, $g_{11\dots 1} = -g_{22\dots 2} = g$, all other components vanish, here g is the coupling constant. As shown in Fig. 2, we change the orientation of incoming momentum r in Fig. 2(a) as outgoing line in Fig. 2(b) with energy-momentum conservation $p + q + \dots - r = 0$. After absorbing $Q^\dagger$ associated with outgoing momentum r and all Q associated with incoming momentum into the vertex, this new bare vertex in Fig. 2(b) should be

$$\gamma_{\beta\beta'\dots\beta''}(p, q, \dots -r) = Q_{\beta a} Q_{\beta' b} \dots Q_{\beta'' c}^\dagger g_{ab\dots c}. \quad (14)$$

From Eq.(11) and (12) we know $Q_{\beta'' c}^\dagger = Q_{c\beta''}^\dagger$. Insert it into Eq.(14) and then compare to Eq.(13), we see clearly that the bare vertex $\gamma_{\beta\beta'\dots\beta''}(p, q, \dots r)$ with $p + q + \dots + r = 0$ is the same as bare vertex $\gamma_{\beta\beta'\dots\beta''}(p, q, \dots -r)$ with $p + q + \dots - r = 0$. This property indicates that the new bare vertex defined here is independent of the orientation of the momentum. In the following we can drop the momentum arguments for the bare vertex. Substituting $Q_{a1}, Q_{a2}, Q_{r1}, Q_{r2}$ in Eq.(11) into Eq.(13), we can express the n-point new bare vertex in the (r, a) basis as

$$\gamma_{\alpha_1\alpha_2\dots\alpha_n} = \left(\frac{1}{\sqrt{2}}\right)^n g [1 - (-1)^{n_a(\alpha_1, \alpha_2, \dots, \alpha_n)}], \quad (15)$$

where $n_a(\alpha_1, \alpha_2, \dots, \alpha_n)$ is the number of a indices among $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$. As this number is even, the bare vertex is zero, so that in the (r, a) basis there are a half of bare vertices vanish and other are only related to the coupling constant and independent of the thermal distribution function. This property will help us to simplify the calculation greatly. For example, the explicit form of three-point bare vertices read

$$\begin{aligned} \gamma_{aaa} &= \gamma_{arr} = \gamma_{rra} = \gamma_{rar} = \frac{g}{\sqrt{2}}, \\ \gamma_{aar} &= \gamma_{ara} = \gamma_{rrr} = \gamma_{raa} = 0. \end{aligned} \quad (16)$$




FIG. 1. The new bare vertex and propagator $\bar{G}_{\alpha\beta}$ after absorbing $Q^\dagger$ and Q into the left and right vertex in our deduced new Feynman rule.




FIG. 2. The bare vertex in the (r, a) basis. The difference between (a) and (b) is that the orientation of momentum r is changed.




FIG. 3. The one-loop Feynman diagram for calculating the retarded and advanced self-energy in ϕ^3 theory.

As first application we use this new Feynman rule to calculate the retarded (or advanced) self-energy in ϕ^3 theory. As shown in Fig. 3, the self-energy in the (r, a) basis can be expressed as

$$-i\Sigma_{\beta\beta'}(q) = \int \frac{d^4p}{(2\pi)^4} (-i\gamma_{\alpha\beta\delta}) (i\Delta_{\alpha\alpha'}(p)) (-i\gamma_{\alpha'\beta'\delta'}) (\Delta_{\delta'\delta}(r)), \quad (17)$$

where $r = p + q$. Noticing propagator $\Delta_{aa} = 0$, the poles of $\Delta_{ra}(p)\Delta_{ra}(r)$ and $\Delta_{ar}(p)\Delta_{ar}(r)$ are both on the same side of the real axis in the complex p^0 plane, we have

$$\int_{-\infty}^{+\infty} \frac{dp^0}{2\pi} \Delta_{ra}(p)\Delta_{ra}(r) = \int_{-\infty}^{+\infty} \frac{dp^0}{2\pi} \Delta_{ar}(p)\Delta_{ar}(r) = 0. \quad (18)$$

Then four components of the self-energy can be derived as

$$-i\Sigma_{ra}(q) = g^2 \int \frac{d^4p}{(2\pi)^4} \{n(r_0)\Delta_{ar}(p)[\Delta_{ra}(r) - \Delta_{ar}(r)] + n(p_0)\Delta_{ra}(r)[\Delta_{ra}(p) - \Delta_{ar}(p)]\}, \quad (19)$$

$$-i\Sigma_{ar}(q) = g^2 \int \frac{d^4p}{(2\pi)^4} \{n(p_0)\Delta_{ar}(r)[\Delta_{ra}(p) - \Delta_{ar}(p)] + n(r_0)\Delta_{ra}(p)[\Delta_{ra}(r) - \Delta_{ar}(r)]\}, \quad (20)$$

$$-i\Sigma_{aa}(q) = [2n(q_0) + 1][\Sigma_{ra}(q) - \Sigma_{ar}(q)], \quad (21)$$

$$-i\Sigma_{rr}(q) = 0. \quad (22)$$

Here $n(k_0) = 1/[\exp(\beta k_0) - 1]$ is the Bose thermal distribution. It is easy to show that $\Sigma_{ar}(q)$ and $\Sigma_{ra}(q)$ are the retarded and advanced self-energy which correspond to the retarded and advanced analytical continuation of the self-energy calculated in the ITF [12]. Eq.(21) corresponds to the Fluctuation-Dissipation theorem in linear response theory [13].

The one-loop three-point vertex correction is illustrated in Fig. 4, the general expression for this vertex correction in the (r, a) basis can be expressed as

$$-i\Gamma_{\alpha\beta\delta}(p, q, r) = \int \frac{d^4l_1}{(2\pi)^4} (-i\gamma_{\alpha\alpha'}\alpha'') [i\Delta_{\alpha'\beta''}(l_1)] (-i\gamma_{\beta\beta'}\beta'') [i\Delta_{\beta'\delta''}(l_2)] (-i\gamma_{\delta\delta'}\delta'') [i\Delta_{\delta'\alpha''}(l_3)], \quad (23)$$

where $p + q + r = 0$ and $l_1 + l_2 + l_3 = 0$.




FIG. 4. The Feynman diagram for calculating one-loop vertex correction in the (r, a) basis in ϕ^3 theory.

Using $\Delta_{aa} = 0$ and the l_1^0 -integral over terms of the type $\Delta_{ra}(l_1)\Delta_{ra}(l_2)\Delta_{ra}(l_3)$ and $\Delta_{ar}(l_1)\Delta_{ar}(l_2)\Delta_{ar}(l_3)$ vanish because the poles of these two terms are both on the same side of the real axis in the complex l_1^0 plane, we deduce eight components of one-loop vertex correction in ϕ^3 theory as

$$-i\Gamma_{rra}(p, q, r) = \frac{2g^3}{(\sqrt{2})^3} \int \frac{d^4l_1}{(2\pi)^4} \left\{ n(l_{01})[\Delta_{ra}(l_1) - \Delta_{ar}(l_1)]\Delta_{ar}(l_2)\Delta_{ra}(l_3) \right.$$

$$+ n(l_{02})[\Delta_{ra}(l_2) - \Delta_{ar}(l_2)]\Delta_{ra}(l_1)\Delta_{ra}(l_3) + n(l_{03})[\Delta_{ra}(l_3) - \Delta_{ar}(l_3)]\Delta_{ar}(l_1)\Delta_{ar}(l_2) \left. \right\}, \quad (24)$$

$$-i\Gamma_{rar}(p, q, r) = \frac{2g^3}{(\sqrt{2})^3} \int \frac{d^4l_1}{(2\pi)^4} \left\{ n(l_{01})[\Delta_{ra}(l_1) - \Delta_{ar}(l_1)]\Delta_{ra}(l_2)\Delta_{ra}(l_3) + n(l_{02})[\Delta_{ra}(l_2) - \Delta_{ar}(l_2)]\Delta_{ar}(l_1)\Delta_{ar}(l_3) + n(l_{03})[\Delta_{ra}(l_3) - \Delta_{ar}(l_3)]\Delta_{ar}(l_1)\Delta_{ra}(l_2) \right\}, \quad (25)$$

$$-i\Gamma_{arr}(p, q, r) = \frac{2g^3}{(\sqrt{2})^3} \int \frac{d^4l_1}{(2\pi)^4} \left\{ n(l_{01})[\Delta_{ra}(l_1) - \Delta_{ar}(l_1)]\Delta_{ar}(l_2)\Delta_{ar}(l_3) + n(l_{02})[\Delta_{ra}(l_2) - \Delta_{ar}(l_2)]\Delta_{ra}(l_1)\Delta_{ar}(l_3) + n(l_{03})[\Delta_{ra}(l_3) - \Delta_{ar}(l_3)]\Delta_{ra}(l_1)\Delta_{ra}(l_2) \right\}, \quad (26)$$

$$-i\Gamma_{aar}(p, q, r) = \frac{g^3}{(\sqrt{2})^3} \int \frac{d^4l_1}{(2\pi)^4} \left\{ [\Delta_{ra}(l_1) - \Delta_{ar}(l_1)]\Delta_{ra}(l_2)\Delta_{ar}(l_3)[N_2 - N_3]N_1 + [\Delta_{ra}(l_1) - \Delta_{ar}(l_1)]\Delta_{ar}(l_2)\Delta_{ar}(l_3)[1 - N_1N_2] + [\Delta_{ra}(l_1) - \Delta_{ar}(l_1)]\Delta_{ra}(l_2)\Delta_{ra}(l_3)[N_1N_3 - 1] \right\}, \quad (27)$$

$$-i\Gamma_{raa}(p, q, r) = \frac{g^3}{(\sqrt{2})^3} \int \frac{d^4l_1}{(2\pi)^4} \left\{ [\Delta_{ra}(l_2) - \Delta_{ar}(l_2)]\Delta_{ra}(l_1)\Delta_{ra}(l_3)[N_1N_2 - 1] + [\Delta_{ra}(l_2) - \Delta_{ar}(l_2)]\Delta_{ar}(l_1)\Delta_{ra}(l_3)[N_2N_3 - N_2N_1] + [\Delta_{ra}(l_2) - \Delta_{ar}(l_2)]\Delta_{ar}(l_1)\Delta_{ar}(l_3)[1 - N_2N_3] \right\}, \quad (28)$$

$$-i\Gamma_{ara}(p, q, r) = \frac{g^3}{(\sqrt{2})^3} \int \frac{d^4l_1}{(2\pi)^4} \left\{ [\Delta_{ra}(l_3) - \Delta_{ar}(l_3)]\Delta_{ar}(l_1)\Delta_{ar}(l_2)[1 - N_1N_3] + [\Delta_{ra}(l_3) - \Delta_{ar}(l_3)]\Delta_{ra}(l_1)\Delta_{ar}(l_2)[N_1N_3 - N_2N_3] + [\Delta_{ra}(l_3) - \Delta_{ar}(l_3)]\Delta_{ra}(l_1)\Delta_{ra}(l_2)[N_2N_3 - 1] \right\}, \quad (29)$$

$$-i\Gamma_{aaa}(p, q, r) = \frac{g^3}{(\sqrt{2})^3} \int \frac{d^4l_1}{(2\pi)^4} \left\{ N_1N_2N_3 \times [\Delta_{ra}(l_1)\Delta_{ra}(l_2)\Delta_{ra}(l_3) - \Delta_{ar}(l_1)\Delta_{ar}(l_2)\Delta_{ar}(l_3)] + (N_1 - N_1N_2N_3)[\Delta_{ra}(l_1) - \Delta_{ar}(l_1)]\Delta_{ra}(l_2)\Delta_{ar}(l_3) + (N_2 - N_1N_2N_3)[\Delta_{ra}(l_2) - \Delta_{ar}(l_2)]\Delta_{ar}(l_1)\Delta_{ar}(l_3) + (N_3 - N_1N_2N_3)[\Delta_{ra}(l_3) - \Delta_{ar}(l_3)] \times \Delta_{ra}(l_1)\Delta_{ar}(l_2) \right\}, \quad (30)$$

$$-i\Gamma_{rrr}(p, q, r) = 0. \quad (31)$$

Here $N_i = 1 + 2n(l_{0i})$. Eq.(22) and (31) indicate the two-point self-energy and three-point vertex with all indices being r vanish, this property is also true for n -point vertex in $n > 3$ case [10].

Denote $N_{k^0} = 1 + 2n(k^0)$ for $k^0 = p^0, q^0, r^0$. As $p + q + r = 0$ and $l_1 + l_2 + l_3 = 0$, we have

$$N_{p^0}[N_3 - N_1] = N_3N_1 - 1, \quad (32)$$

$$N_{q^0}[N_1 - N_2] = N_1N_2 - 1, \quad (33)$$

$$N_{r^0}[N_2 - N_3] = N_2N_3 - 1. \quad (34)$$

Using above equations we can verify that three-point vertex corrections in Eq.(24)-(31) satisfy following relations:

$$\Gamma_{ara} = N_{p^0}(\Gamma_{rar}^* - \Gamma_{rra}) + N_{r^0}(\Gamma_{rar}^* - \Gamma_{arr}), \quad (35)$$

$$\Gamma_{raa} = N_{q^0}(\Gamma_{arr}^* - \Gamma_{rra}) + N_{r^0}(\Gamma_{arr}^* - \Gamma_{rar}), \quad (36)$$

$$\Gamma_{aar} = N_{p^0}(\Gamma_{rra}^* - \Gamma_{rar}) + N_{q^0}(\Gamma_{rra}^* - \Gamma_{arr}), \quad (37)$$

$$\begin{aligned} \Gamma_{aaa} &= \Gamma_{arr}^* + \Gamma_{rar}^* + \Gamma_{rra}^* + N_{q^0}N_{r^0}(\Gamma_{arr} + \Gamma_{arr}^*) \\ &\quad + N_{p^0}N_{r^0}(\Gamma_{rar} + \Gamma_{rar}^*) \\ &\quad + N_{p^0}N_{q^0}(\Gamma_{rra} + \Gamma_{rra}^*). \end{aligned} \quad (38)$$

These relations are just the generalized Fluctuation-Dissipation theorem for three-point nolinear response function which is in agreement with the result obtained in our previous work [10]. Eq.(35)-(38) together with Eq.(31) indicate that, among these vertex functions $\{\Gamma_{rra}, \Gamma_{rar}, \Gamma_{arr}, \Gamma_{raa}, \Gamma_{ara}, \Gamma_{aar}, \Gamma_{aaa}, \Gamma_{rrr}\}$ only three components are independent.

In Ref. [9] Aurenche and Becherrawy developed a new Feynman rule which denoted by R, A to calculate retarded and advanced vertex function. The main difference between their and our Feynman rule is following: In the (R, A) Feynman rule the matrix propagator is diagonal and independent of the thermal distribution function; the thermal distribution function is absorbed into the vertex, the defined new vertex depends on the relative orientation of the momenta and only two of the vertex components vanish. In our (r, a) Feynman rule the matrix propagator is non-diagonal, one component is zero and another component Δ_{rr} is related to thermal distribution function; the defined new bare vertex is independent of thermal distribution function and only related to the coupling constant; this new bare vertex doesn't rely on the orientation of the momenta and one half of the bare vertex components vanish. So that in the practical calculation both (R, A) and (r, a) Feynman rule have their own advantage and disadvantage. We should point out that, in our (r, a) basis the Green function is defined with time-ordering and have explicit physical interpretation (see Eq.(4)-(7) and Ref. [10]), it works in both coordinate and momentum space; but the (R, A) Feynman rule, as stated by the authors themselves in Ref. [9], only works in the momentum space, and the (R, A) Green function have no explicit reference to a definite time-ordering.

By suitable combination we can get the (R, A) vertex from the (r, a) vertex. For three-point vertex the relation between them can be expressed as

$$\tilde{\Gamma}_{RRR} = \tilde{\Gamma}_{AAA} = 0, \quad (39)$$

$$\tilde{\Gamma}_{RRA} = \Gamma_{rra}, \quad (40)$$

$$\tilde{\Gamma}_{RAR} = \Gamma_{rar}, \quad (41)$$

$$\tilde{\Gamma}_{ARR} = \Gamma_{arr}, \quad (42)$$

$$\tilde{\Gamma}_{AAR} = -\frac{1}{2}[N_{p^0} + N_{q^0}]\Gamma_{rra}^*, \quad (43)$$

$$\tilde{\Gamma}_{ARA} = -\frac{1}{2}[N_{p^0} + N_{r^0}]\Gamma_{rar}^*, \quad (44)$$

$$\tilde{\Gamma}_{RAA} = -\frac{1}{2}[N_{p^0} + N_{r^0}]\Gamma_{arr}^*. \quad (45)$$

In summary, the new Feynman rule in the (r, a) basis is deduced from the Keldysh transformation in the CTPF. This Feynman rule can be used to calculate the retarded and advanced Green function directly. In this new Feynman rule, the bare vertex is only related with the coupling constant and independent of the orientation of the momenta, one half of bare vertex components vanish; the new propagator depends on retarded, advanced propagator and the thermal distribution function. As application, the one-loop self-energy and three-point vertex correction in ϕ^3 theory are calculated in the (r, a) basis, and the Fluctuation-Dissipation theorem is verified in the two-point and three-point cases. The difference between our (r, a) Feynman rule and the (R, A) Feynman rule introduced by Aurenche and Becherrawy [9] is discussed.

ACKNOWLEDGMENTS

This work was supported by the National Natural Science Foundation of China (NSFC) under Grant Nos. 10135030 and 19928511.

-
- [1] Schwinger J (1961) *J. Math. Phys.* **2** 407
 - [2] Keldysh L V (1965) *Sov. Phys. JETP* **20** 1018
 - [3] Chou K C, Su Z B, Hao B L and Yu L (1985) *Phys. Rep.* **118** 1
 - [4] Wang E K, Heinz U (1999) *Phys. Lett. B* **471** 208; and hep-th/0201116
 - [5] Wang E K, Heinz U and Zhang X F (1996) *Phys. Rev. D* **53** 5978
 - [6] Hou D F, Wang E K and Heinz U (1998) *J. Phys. G* **24** 1861
 - [7] Aarts G and Smit J (1998) *Nucl. Phys. B* **511** 451
 - [8] van Eijck M A, Kobes R and van Weert C G (1994) *Phys. Rev. D* **50** 4097
 - [9] Aurenche P and Becherrawy T (1992) *Nucl. Phys. B* **379** 259
 - [10] Wang E K and Heinz U, (2002) *Phys. Rev. D* (in press), hep-th/9809016
 - [11] Lehmann H, Symanzik K and Zimmermann W (1957) *Nuovo Cimento* **6** 319
 - [12] Weldon H A (1983) *Phys. Rev. D* **28** 2007
 - [13] Callen H B and Welton T A (1951) *Phys. Rev.* **83** 34